

University of Groningen

Study of the isospin character of 1-states using hadronic probes at intermediate energies

Derya, V.; Endres, J.; Harakeh, M.N.; Savran, D.; Spieker, M.; Wörtche, H.J.; Zilges, A.

Published in:
EPJ Web of Conferences

DOI:
[10.1051/epjconf/20159301041](https://doi.org/10.1051/epjconf/20159301041)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Derya, V., Endres, J., Harakeh, M. N., Savran, D., Spieker, M., Wörtche, H. J., & Zilges, A. (2015). Study of the isospin character of 1-states using hadronic probes at intermediate energies. *EPJ Web of Conferences*, 93, [01041]. <https://doi.org/10.1051/epjconf/20159301041>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Study of the isospin character of 1^- states using hadronic probes at intermediate energies

V. Derya^{1,a}, J. Endres¹, M.N. Harakeh^{2,3}, D. Savran^{4,5}, M. Spieker¹, H.J. Wörtche², and A. Zilges¹

¹*Institut für Kernphysik, Universität zu Köln, 50937 Köln, Germany*

²*Kernfysisch Versneller Instituut, University of Groningen, 9747 AA Groningen, The Netherlands*

³*GANIL, CEA/DSM-CNRS/IN2P3, 14076 Caen, France*

⁴*ExtreMe Matter Institute EMMI and Research Division, GSI, 64291 Darmstadt, Germany*

⁵*Frankfurt Institute for Advanced Studies FIAS, 60438 Frankfurt am Main, Germany*

Abstract. The complementary (γ, γ') and $(\alpha, \alpha'\gamma)$ reactions were used to study the isospin properties of low-lying $E1$ excitations in the doubly-magic nucleus ^{48}Ca . In contrast to heavier nuclei, a state-to-state change in isospin character was revealed in ^{48}Ca and a dominant isoscalar excitation was found which is interpreted as an isoscalar oscillation. Recently, protons at 80 MeV were used as an additional hadronic probe in a p - γ coincidence experiment on ^{140}Ce for the first time. Results of the experiments on ^{48}Ca and first results of the ^{140}Ce will be presented in this contribution.

1 Introduction

Low-lying electric dipole excitations, in particular the electric Pygmy Dipole Resonance (PDR), were investigated in neutron-rich nuclei using various experimental methods [1]. Aiming at isospin characters of these 1^- states, it has been demonstrated that the complementary studies of real-photon scattering and high-resolution $(\alpha, \alpha'\gamma)$ coincidence experiments at 136 MeV allow separating isovector and isoscalar dipole response which is important for a deeper understanding of the underlying mechanisms generating the dipole strength [2]. Systematic studies using these complementary probes revealed a splitting into low-lying isospin-mixed $E1$ excitations and higher-lying dominantly isovector $E1$ excitations in neutron-magic, proton-magic, and non-magic nuclei (namely ^{140}Ce , ^{138}Ba , ^{124}Sn , and ^{94}Mo) [2–5]. This experimentally observed common feature of the low-lying $E1$ response is reproduced by several theoretical calculations (see, e.g., Refs. [6–9]) which suggest a distinction between a low-lying neutron-skin oscillation mode and a transitional mode towards the well-known isovector Giant Dipole Resonance (IVGDR) [10]. Recently, the crossover between the neutron-skin mode and the higher-lying proton-neutron oscillation mode was investigated in detail by means of random-phase approximation (RPA) calculations [11] and a decomposition method was introduced which might be important for a robust comparison with experimental data. In addition to the aforementioned α -particles, the hadronic probe of ^{17}O at 20 MeV/u was recently used in a high-resolution experiment on ^{208}Pb at Legnaro National Laboratory [12].

In the following, we present results of two experiments using the hadronic probes of α particles at 136 MeV and protons at 80 MeV which we performed to achieve a better understanding of the nature of low-lying $E1$ excitations in ^{48}Ca and ^{140}Ce . The $^{48}\text{Ca}(\alpha, \alpha'\gamma)$ and $^{140}\text{Ce}(p, p'\gamma)$ experiments go beyond the systematic study of low-lying dipole excitations in $(\alpha, \alpha'\gamma)$ and (γ, γ') experiments in two ways: On the one hand, by studying a much lighter neutron-rich nucleus and, on the other hand, by using a complementary hadronic probe with a higher energy per nucleon, namely protons at 80 MeV which penetrate more deeply into the nucleus.

2 Experimental setup and results

Both experiments were performed at the KVI Groningen, The Netherlands, exploiting the Big-Bite Spectrometer in combination with an array of HPGe detectors for γ spectroscopy. The scattered particles and the de-exciting γ -rays were acquired in hardware coincidence. In the data-analysis process, particle- γ coincidence matrices are constructed which are then used to generate energy spectra with specific energy conditions on the excitation energy (E_x) and the γ -ray energy (E_γ). Gates on ground-state transitions ($E_x = E_\gamma$) highly increase the sensitivity to dipole transitions which dominantly decay via this channel. Details on the experimental setup and data analysis tools can be found in Ref. [13]. Results of the two experiments on ^{48}Ca and ^{140}Ce are presented in the following subsections. Main experimental parameters are given in Table 2.


^ae-mail: derya@ikp.uni-koeln.de

Table 1. Experimental parameters for the $^{48}\text{Ca}(\alpha, \alpha'\gamma)$ and $^{140}\text{Ce}(p, p'\gamma)$ experiments.

	$^{48}\text{Ca}(\alpha, \alpha'\gamma)$	$^{140}\text{Ce}(p, p'\gamma)$
target thickness	1.7 mg/cm ²	20 mg/cm ²
isotopic enrichment	99%	99.72%
beam energy	136 MeV	80 MeV
average beam current	0.9 pA	0.6 pA
BBS central angle	5.8°	5.6°
collected charge	794 μC	408 μC


2.1 Results of the $^{48}\text{Ca}(\alpha, \alpha'\gamma)$ experiment

Low-lying $E1$ excitations and their $B(E1)$ strength distributions in stable Ca isotopes, including the doubly-magic ^{48}Ca , have been studied in several (γ, γ') experiments [14, 15]. Hence, the results of the $^{48}\text{Ca}(\alpha, \alpha'\gamma)$ experiment of this work, allow a comparison of singles α -scattering cross sections and reduced $B(E1)^\uparrow$ transition strengths [14] (presented in Fig. 1) which gives access to the isospin character of low-lying 1^- states in ^{48}Ca . It reveals a state-to-state dependent isospin character of the 1^- states where isoscalar, isospin-mixed, and isovector dipole excitations are in close vicinity. An isospin splitting as observed in the heavier nuclei is not present in ^{48}Ca . Remarkable is also a strong dominantly isoscalar state at 7.6 MeV close to a strong isovector state at 7.3 MeV. The strong isoscalar state is interpreted as a pure isoscalar oscillation on basis of RPA calculations [16] under consideration of the corresponding velocity fields and transition densities. Furthermore, the well-separated dominant isovector and isoscalar states at 7.3 MeV and 7.6 MeV, respectively, are an optimal test case for the investigation of isospin mixing in a two-state mixing approach. A value of 0.061(6) was determined for the squared mixing amplitude. This results in an isospin-mixing matrix element of 85(3) keV. Further results of this experiment were presented in more detail in Ref. [17].


**Figure 1.** Cross sections obtained in the $^{48}\text{Ca}(\alpha, \alpha'\gamma)$ experiment (upper panel) in comparison with results of a real-photon scattering experiment [14] (lower panel).

2.2 Results of the $^{140}\text{Ce}(p, p'\gamma)$ experiment

The neutron-magic ^{140}Ce was the first nucleus for which an isospin splitting of low-lying $E1$ excitations was observed and it is the first nucleus for which the studies were extended by a $(p, p'\gamma)$ experiment at intermediate energy. The summed γ -ray spectrum of all HPGe detectors with a gate on ground-state transitions obtained in the $^{140}\text{Ce}(p, p'\gamma)$ experiment is shown in Fig. 2.

**Figure 2.** Summed γ -ray spectrum of all HPGe detectors with gate on $E_x = E_\gamma$ measured in the $^{140}\text{Ce}(p, p'\gamma)$ experiment. Dipole transitions are marked with stars.

Dipole transitions were identified via their ground-state transition energy in comparison with (γ, γ') data [18, 19]. Singles proton-scattering cross sections were determined on basis of this spectrum. Compared to the results from the inelastic α -scattering experiment, the proton-scattering cross sections are almost one order of magnitude smaller. The ratio of the cross sections is shown in Fig. 2. It is nearly constant with an average value of $(d\sigma/d\Omega_\alpha)/(d\sigma/d\Omega_p) = 6.2(7)$. The central part of the nucleon-nucleon interaction is strongly energy-dependent [20] and decreases with increasing energy from 34 MeV/u to 80 MeV/u. This also leads to a higher degree of transparency for the proton probe and additional cancellation effects for dipole transitions. The experimentally determined ratio for the cross sections is reproduced by Distorted-Wave Born Approximation (DWBA) calculations.

**Figure 3.** Ratios of singles α -scattering cross sections [2] and proton-scattering cross sections (this work) for the dipole excitations observed in both experiments. The horizontal line indicates the average ratio.

3 Summary and outlook

Inelastic scattering experiments using hadronic probes at intermediate energies, namely an $(\alpha, \alpha'\gamma)$ experiment at $E_\alpha = 136$ MeV on the doubly-magic ^{48}Ca and a $(p, p'\gamma)$ experiment at $E_p = 80$ MeV on the neutron-magic ^{140}Ce were performed at KVI Groningen to further study the nature of low-lying $E1$ strength. These experiments complement previous real-photon scattering experiments on both nuclei [14, 18] and an additional $^{140}\text{Ce}(\alpha, \alpha'\gamma)$ experiment [2]. In contrast to the observations in heavier nuclei, low-lying $E1$ excitations in the lighter nucleus ^{48}Ca show a state-to-state dependent isospin character. Furthermore, a strong isoscalar dipole state was revealed which supports theoretical predictions of a strong isoscalar oscillation [16].

For the dipole excitations in ^{140}Ce , the singles proton-scattering cross sections are considerably smaller compared to the α -scattering cross sections. Nevertheless, the general excitation pattern seems to be very similar which is supported by a nearly constant ratio of these cross sections.

In the future, different experimental techniques will be combined to achieve a deeper understanding of co-existing electric dipole modes throughout the nuclear landscape. Essential observables testing the underlying structures more thoroughly include γ -decay branchings, isospin characters, as well as single-particle contents. In particular, the isospin character of 1^- states will be addressed in particle- γ coincidence experiments with hadronic probes at intermediate energies at iThemba LABS in Somerset West, South Africa, and within an experimental PDR campaign with the CAGRA Clover-detector array at RCNP in Osaka, Japan.

Acknowledgment

This work was supported by the DFG (ZI 510/4-2), by the European Commission within the Sixth Framework Programme through I3-EURONS (contract No. RII3-CT-2004-506065), by the Alliance Program of the Helmholtz Association (HA216/EMMI), and by the Helmholtz International Center for FAIR (HIC for FAIR).

References

- [1] D. Savran, T. Aumann, A. Zilges, Prog. Part. Nucl. Phys. **70**, 210 (2013)
- [2] D. Savran, M. Babilon, A.M. van den Berg, M.N. Harakeh, J. Hasper, A. Matic, H.J. Wörtche, A. Zilges, Phys. Rev. Lett. **97**, 172502 (2006)
- [3] J. Endres, D. Savran, A.M. van den Berg, P. Dendooven, M. Fritzsche, M.N. Harakeh, J. Hasper, H.J. Wörtche, A. Zilges, Phys. Rev. C **80**, 034302 (2009)
- [4] J. Endres, E. Litvinova, D. Savran, P.A. Butler, M.N. Harakeh, S. Harissopulos, R.D. Herzberg, R. Krücken, A. Lagoyannis, N. Pietralla et al., Phys. Rev. Lett. **105**, 212503 (2010)
- [5] V. Derya, J. Endres, M. Elvers, M.N. Harakeh, N. Pietralla, C. Romig, D. Savran, M. Scheck, F. Siebenhühner, V.I. Stoica et al., Nucl. Phys. A **906**, 94 (2013)
- [6] N. Tsoneva, H. Lenske, Phys. Rev. C **77**, 024321 (2008)
- [7] N. Paar, Y.F. Niu, D. Vretenar, J. Meng, Phys. Rev. Lett. **103**, 032502 (2009)
- [8] E.G. Lanza, A. Vitturi, E. Litvinova, D. Savran, Phys. Rev. C **89**, 041601 (2014)
- [9] E. Litvinova, P. Ring, V. Tselyaev, K. Langanke, Phys. Rev. C **79**, 054312 (2009)
- [10] M.N. Harakeh, A. van der Woude, *Giant Resonances* (Oxford University Press, New York, 2001)
- [11] H. Nakada, T. Inakura, H. Sawai, Phys. Rev. C **87**, 034302 (2013)
- [12] F.C.L. Crespi, A. Bracco, R. Nicolini, D. Mengoni, L. Pellegrini, E.G. Lanza, S. Leoni, A. Maj, M. Kmiecik, R. Avigo et al., Phys. Rev. Lett. **113**, 012501 (2014)
- [13] D. Savran, A.M. van den Berg, M.N. Harakeh, K. Ramspeck, H.J. Wörtche, A. Zilges, Nucl. Instr. and Meth. A **564**, 267 (2006)
- [14] T. Hartmann, J. Enders, P. Mohr, K. Vogt, S. Volz, A. Zilges, Phys. Rev. C **65**, 034301 (2002)
- [15] J. Isaak, D. Savran, M. Fritzsche, D. Galaviz, T. Hartmann, S. Kamedzhiev, J.H. Kelley, E. Kwan, N. Pietralla, C. Romig et al., Phys. Rev. C **83**, 034304 (2011)
- [16] P. Papakonstantinou, H. Hergert, V.Yu. Ponomarev, R. Roth, Phys. Lett. B **709**, 270 (2012)
- [17] V. Derya, D. Savran, J. Endres, M.N. Harakeh, H. Hergert, J.H. Kelley, P. Papakonstantinou, N. Pietralla, V.Yu. Ponomarev, R. Roth et al., Phys. Lett. B **730**, 288 (2014)
- [18] S. Volz, N. Tsoneva, M. Babilon, M. Elvers, J. Hasper, R.D. Herzberg, H. Lenske, K. Lindenberg, D. Savran, A. Zilges, Nucl. Phys. A **779**, 1 (2006)
- [19] B. Löher (2014), private communication
- [20] W.G. Love, M.A. Franey, Phys. Rev. C **24**, 1073 (1981)

